

CLASS :VIII

SUBJECT :SCIENCE

LESSON :LIGHT

PREPARED BY:

K.PRAKASH .,M.Sc., M.A., M.Ed., M.Phil.,

BRINDAVAN MATRIC HR SEC SCHOOL,ATHUR


CHENGALPATTU DISTRICT.

PH:8489006155

LIGHT

TYPES OF MIRRORS

- **We use mirrors in our daily life for various purposes.**
- **We use them for decoration.**
- **In vehicles, they are used as rear view mirrors.**
- **They are also used in scientific apparatus, like telescope.**
- **The mirror is an optical device with a polished surface that reflects the light falling on it.**
- **A typical mirror is a glass sheet coated with aluminium or silver on one of its sides to produce an image.**
- **Mirrors have a plane or curved surface.**
- **Curved mirrors have surfaces that are spherical, cylindrical, parabolic and ellipsoid.**
- **The shape of a mirror determines the type of image it forms.**
- **Plane mirrors form the perfect image of an object.**
- **Whereas, curved mirrors produce images that are either enlarged or diminished.**
- **You would have studied about plane mirrors in your lower classes.**
- **In this section, you will study about spherical and parabolic mirrors.**



Method of coating a glass plate with a thin layer of reflecting metals was in practice during the 16th century in Venice, Italy. They used an amalgam of tin and mercury for this purpose. Nowadays, a thin layer of molten aluminium or silver is used for coating glass plates that will then become mirrors.


SPHERICAL MIRRORS

- Spherical mirrors are one form of curved mirrors.
- If the curved mirror is a part of a sphere, then it is called a 'spherical mirror'.
- It resembles the shape of a piece cut out from a spherical surface.
- One side of this mirror is silvered and the reflection of light occurs at the other side.

CONCAVE MIRRORS

- A spherical mirror, in which the reflection of light occurs at its concave surface, is called a concave mirror.
- These mirrors magnify the object placed close to them.
- The most common example of a concave mirror is the make-up mirror.


CONCAVE MIRRORS



Convex mirror

- Convex mirror A spherical mirror, in which the reflection of light occurs at its convex surface, is called a convex mirror.
- The image formed by these mirrors is smaller than the object.
- Most common convex mirrors are rear viewing mirrors used in vehicles.

Convex mirror



Convex mirror




Convex mirrors used in vehicles as rear-view mirrors are labeled with the safety warning: 'Objects in the mirror are closer than they appear' to warn the drivers. This is because inside the mirrors, vehicles will appear to be coming at a long distance.

Parabolic mirrors

- A parabolic mirror is one type of curved mirror, which is in the shape of a parabola.
- It has a concave reflecting surface and this surface directs the entire incident beam of light to converge at its focal point.
- In the same way, light rays generated by the source placed at this focal point will fall on this surface and they will be diverged in a direction, which is parallel to the principal axis of the parabolic mirror.
- Hence, the light rays will be reflected to travel a long distance, without getting diminished.
- Parabolic mirrors, also known as parabolic reflectors, are used to collect or project energy such as light, heat, sound and radio waves.
- They are used in reflecting telescopes, radio telescopes and parabolic microphones. They are also used in solar cookers and solar water heaters.

PARABOLIC MIRRORS



A Parabolic Mirror



Figure 3.3 Parabolic mirror



The principle behind the working of a parabolic mirror has been known since the Greco-Roman times. The first mention of these structures was found in the book, 'On Burning Mirrors', written by the mathematician Diocles. They were also studied in the 10th century, by a physicist called Ibn Sahl. The first parabolic mirrors were constructed by Heinrich Hertz, a German physicist, in the form of reflector antennae in the year 1888.

TERMS RELATED TO SPHERICAL MIRRORS

- In order to understand the image formation in spherical mirrors, you need to know about some of the terms related to them.
- **Center of Curvature:** It is the center of the sphere from which the mirror is made. It is denoted by the letter C in the ray diagrams.
- (A ray diagram represents the formation of an image by the spherical mirror. You will study about them in your next class).
- **Pole:** It is the geometric centre of the spherical mirror. It is denoted by the letter P.
- **Radius of Curvature:** It is the distance between the center of the sphere and the vertex. It is shown by the letter R in ray diagrams. (The vertex is the point on the mirror's surface where the principal axis meets the mirror. It is also called as 'pole'.)

TERMS RELATED TO SPHERICAL MIRRORS

- **Principal Axis:** The line joining the pole of the mirror and its center of curvature is called principal axis.
- **Focus:** When a beam of light is incident on a spherical mirror, the reflected rays converge (concave mirror) at or appear to diverge from (convex mirror) a point on the principal axis.
- This point is called the 'focus' or 'principal focus'. It is also known as the focal point.
- It is denoted by the letter F in ray diagrams.

TERMS RELATED TO SPHERICAL MIRRORS

Focal length: The distance between the pole and the principal focus is called focal length (f) of a spherical mirror.

There is a relation between the focal length of a spherical mirror and its radius of curvature. The focal length is half of the radius of curvature.

$$\text{That is, focal length} = \frac{\text{Radius of curvature}}{2}.$$



Figure 3.4 Terms related to a spherical mirror

NUMERICALS

PROBLEM :1

The radius of curvature of a spherical mirror is 20 cm. Find its focal length.

Solution:

Radius of curvature = 20 cm

$$\begin{aligned}\text{Focal length (f)} &= \text{Radius of curvature} / 2 \\ &= R/2 \quad = 20 / 2 \quad = 10 \text{ cm}\end{aligned}$$

PROBLEM : 2

Focal length of a spherical mirror is 7 cm. What is its radius of curvature?

Solution:

Focal length = 7 cm

$$\begin{aligned}\text{Radius of curvature (R)} &= 2 \times \text{focal length} \\ &= 2 \times 7 \\ &= 14 \text{ cm}\end{aligned}$$


IMAGES FORMED BY SPHERICAL MIRRORS

- Images formed by spherical mirrors are of two types: i) real image and ii) virtual image.
- Real images can be formed on a screen, while virtual images cannot be formed on a screen.
- Image formed by a convex mirror is always erect, virtual and diminished in size.
- As a result, images formed by these mirrors cannot be projected on a screen.
- The characteristics of an image are determined by the location of the object.
- As the object gets closer to a concave mirror, the image gets larger, until attaining approximately the size of the object, when it reaches the centre of curvature of the mirror.
- As the object moves away, the image diminishes in size and gets gradually closer to the focus, until it is reduced to a point at the focus when the object is at an infinite distance from the mirror.
- The size and nature of the image formed by a convex mirror is given in Table 3.1.

IMAGES FORMED BY CONVEX MIRROR

POSITION OF THE OBJECT	POSITION OF THE IMAGE	IMAGE SIZE	NATURE OF THE IMAGE
At infinity	At F	Highly diminished, point sized	Virtual and erect
Between infinity and the pole (P)	Between P and F	Diminished	Virtual and erect

IMAGES FORMED BY CONVEX MIRROR



IMAGES FORMED BY SPHERICAL MIRRORS







- Concave mirrors form a real image and it can be caught on a screen.
- Unlike convex mirrors, concave mirrors show different image types.
- Depending on the position of the object in front of the mirror, the position, size and nature of the image will vary.

SUMMARY OF IMAGES FORMED BY A CONCAVE MIRROR.

Table 3.2 Image formed by a concave mirror

POSITION OF THE OBJECT	POSITION OF THE IMAGE	IMAGE SIZE	NATURE OF THE IMAGE
At infinity	At F	Highly diminished	Real and inverted
Beyond C	Between C and F	Diminished	Real and inverted
At C	At C	Same size as the object	Real and inverted
Between C and F	Beyond C	Magnified	Real and inverted
At F	At infinity	Highly magnified	Real and inverted
Between F and P	Behind the mirror	Magnified	Virtual and erect

SUMMARY OF IMAGES FORMED BY A CONCAVE MIRROR.



SUMMARY OF IMAGES FORMED BY A CONCAVE MIRROR.

- You can observe from the table that a concave mirror always forms a real and inverted image except when the object is placed between the focus and the pole of the mirror.
- In this position, it forms a virtual and erect image.

APPLICATIONS OF CURVED MIRRORS

❖ CONCAVE MIRRORS:

- 1. Concave mirrors are used while applying make-up or shaving, as they provide a magnified image.
- 2. They are used in torches, search lights and head lights as they direct the light to a long distance.
- 3. They can collect the light from a larger area and focus it into a small spot. Hence, they are used in solar cookers.
- 4. They are used as head mirrors by doctors to examine the eye, ear and throat as they provide a shadow-free illumination of the organ.
- 5. They are also used in reflecting telescopes.

APPLICATIONS OF CONCAVE MIRRORS

❖ CONCAVE MIRRORS:



APPLICATIONS OF CONVEX MIRRORS

❖ CONVEX MIRRORS:

1. Convex mirrors are used in vehicles as rear view mirrors because they give an upright image and provide a wider field of view as they are curved outwards.
2. They are found in the hallways of various buildings including hospitals, hotels, schools and stores. They are usually mounted on a wall or ceiling where hallways make sharp turns.
3. 3. They are also used on roads where there are sharp curves and turns.

APPLICATIONS OF CONVEX MIRRORS



LAWS OF REFLECTION

Activity : 3

- Take a plane mirror and try to focus the light coming from the Sun on a wall.
- Can you see a bright spot on the wall?
- How does it occur?
- It is because the light rays falling on the mirror are bounced onto the wall by it.
- Can you produce the same bright spot with the help of any other object having a rough surface?

LAWS OF REFLECTION

- Not all the objects can produce the same effect as produced by the plane mirror.
- A ray of light, falling on a body having a shiny, polished and smooth surface alone is bounced back.
- This bouncing back of the light rays as they fall on the smooth, shiny and polished surface is called reflection.

LAWS OF REFLECTION


Reflection involves two rays:

i) incident ray and ii) reflected ray.

- The incident ray is the light ray in a medium falling on the shiny surface of a reflecting body.
- After falling on the surface, this ray returns into the same medium. This ray is called the reflected ray.
- An imaginary line perpendicular to the reflecting surface, at the point of incidence of the light ray, is called the normal.
- The relation between the incident ray, the reflected ray and the normal is given as the law of reflection.

LAWS OF REFLECTION

- The incident ray, the reflected ray and the normal at the point of incidence, all lie in the same plane.
- The angle of incidence and the angle of reflection are always equal.



TYPES OF REFLECTION

1.REGULAR REFLECTION 2.IRREGULAR REFLECTION

1.REGULAR REFLECTION:

- When a beam of light (collection of parallel rays) falls on a smooth surface, it gets reflected.
- After reflection, the reflected rays will be parallel to each other.
- Here, the angle of incidence and the angle of reflection of each ray will be equal.
- Hence, the law of reflection is obeyed in this case and thus a clear image is formed.
- This reflection is called 'regular reflection' or 'specular reflection'.
- Example: Reflection of light by a plane mirror and reflection of light from the surface of still water.

REGULAR REFLECTION




Figure 3.8 Regular reflection

IRREGULAR REFLECTION

- In the case of a body having a rough or irregular surface, each region of the surface is inclined at different angles.
- When light falls on such a surface, the light rays are reflected at different angles.
- In this case, the angle of incidence and the angle of reflection of each ray are not equal.
- Hence, the law of reflection is not obeyed in this case and thus the image is not clear.
- Such a reflection is called 'irregular reflection' or 'diffused reflection'.
- Example: Reflection of light from a wall.

IRREGULAR REFLECTION





Figure 3.8 Irregular reflection

MULTIPLE REFLECTION



Activity 4

Take two plane mirrors and keep them perpendicular to each other. Place an object between them. You can see the images of the object. How many images do you see in the mirrors? You can see three images. How is it possible to have three images with two mirrors?



MULTIPLE REFLECTION

- In the activity given above, you observed that for a body kept in between two plane mirrors, which were inclined to each other, you could see many images.
- This is because, the 'image' formed by one mirror acts as an 'object' for the other mirror.
- The image formed by the first mirror acts as an object for the second mirror and the image formed by the second mirror acts as an object for the first mirror.
- Thus, we have three images of a single body. This is known as multiple reflection. This type of reflections can be seen in show rooms and saloons.

MULTIPLE REFLECTION

PROBLEM 3

If two plane mirrors are inclined to each other at an angle of 90° , find the number of images formed.

Solution:

Angle of inclination = 90°


Number of images formed =

$$\frac{360^\circ}{\theta} - 1 = \frac{360^\circ}{90^\circ} - 1 = 4 - 1 = 3$$

KALEIDOSCOPE

- It is a device, which functions on the principle of multiple reflection of light, to produce numerous patterns of images.
- It has two or more mirrors inclined with each other.
- It can be designed from inexpensive materials and the colourful image patterns formed by this will be pleasing to you.
- This instrument is used as a toy for children.

KALEIDOSCOPE



Kaleidoscope



Activity 5

-
- **Take three equal sized plane mirror strips and arrange them in such a way that they form an equilateral triangle.**
- **Cover the sides of the mirrors with a chart paper. With the help of a chart paper cover the bottom of the mirrors also.**
- **Put some coloured things such as pieces of bangles and beads inside it.**
- **Now, cover the top portion with the chart paper and make a hole in it to see.**
- **You can wrap the entire piece with coloured papers to make it attractive.**
- **Now, rotate it and see through its opening. You can see the beautiful patterns.**
- **CAUTION: Be careful while handling the glass pieces. Do this under the supervision of your teacher.**

PERISCOPE

- It is an instrument used for viewing bodies or ships, which are over and around another body or a submarine.
- It is based on the principle of the law of reflection of light.
- It consists of a long outer case and inside this case mirrors or prisms are kept at each end, inclined at an angle of 45° .
- Light coming from the distant body, falls on the mirror at the top end of the periscope and gets reflected vertically downward.
- This light is reflected again by the second mirror kept at the bottom, so as to travel horizontally and reach the eye of the observer.
- In some complex periscopes, optic fibre is used instead of mirrors for obtaining a higher resolution.
- The distance between the mirrors also varies depending on the purpose of using the periscope.

PERISCOPE




Figure 3.11 Periscope used in Submarine

USES OF PERISCOPE

- It is used in warfare and navigation of the submarine.
- In military it is used for pointing and firing guns from a 'bunker'.
- Photographs of important places can be taken through periscopes without trespassing restricted military regions.
- Fibre optic periscopes are used by doctors as endoscopes to view internal organs of the body.

REFRACTION OF LIGHT

- We know that when a light ray falls on a polished surface placed in air, it is reflected into the air itself. When it falls on a transparent material, it is not
- reflected completely, but a part of it is reflected and a part of it is absorbed and most of the light passes through it.
- Through air, light travels with a speed of 3×10^8 m/s, but it cannot travel with the same speed in water or glass, because, optically denser medium such as water and glass offer some resistance to the light rays.
- So, light rays travelling from a rarer medium like air into a denser medium like glass or water are deviated from their straight line path

REFRACTION OF LIGHT

- This bending of light about the normal, at the point of incidence; as it passes from one transparent medium to another is called refraction of light.
- When a light ray travels from the rarer medium into the denser medium, it bends towards the normal and when it travels from the denser medium into the rarer medium, it bends away from the normal.
- You can observe this phenomenon with the help of the activity given below.

REFRACTION OF LIGHT

Activity 6

Take a glass beaker, fill it with water and place a pencil in it. Now, look at the pencil through the beaker. Does it appear straight? No. It will appear to be bent at the surface of the water. Why?



In this activity, the light rays actually travel from the water (a denser medium) into the air (a rarer medium). As you saw earlier, when a light ray travels from a denser medium to a rarer medium, it is deviated from its straight line path. So, the pencil appears to be bent when you see it through the glass of water.

REFRACTIVE INDEX

- Refraction of light in a medium depends on the speed of light in that medium.
- When the speed of light in a medium is more, the bending is less and when the speed of light is less, the bending is more.
- The amount of refraction of light in a medium is denoted by a term known as refractive index of the medium, which is the ratio of the speed of light in the air to the speed of light in that particular medium.
- It is also known as the absolute refractive index and it is denoted by the Greek letter ' μ ' (pronounced as 'mew').
- $\mu = \text{Speed of light in air (c)} / \text{Speed of light in the medium (v)}$

REFRACTIVE INDEX

- Refractive index is a ratio of two similar quantities (speed) and so, it has no unit.
- Since, the speed of light in any medium is less than its speed in air, refractive index of any transparent medium is always greater than 1.

REFRACTIVE INDEX

Table 3.3 Refractive Index of substances

SUBSTANCES	REFRACTIVE INDEX
Air	1.0
Water	1.33
Ether	1.36
Kerosene	1.41
Ordinary Glass	1.5
Quartz	1.56
Diamond	2.41

In general, the refractive index of one medium with respect to another medium is given by the ratio of their absolute refractive indices.

$${}_1\mu_2 = \frac{\text{Absolute refractive index of the second medium}}{\text{Absolute refractive index of the first medium}}$$

$${}_1\mu_2 = \frac{\frac{c}{v_2}}{\frac{c}{v_1}} \quad \text{or} \quad {}_1\mu_2 = \frac{v_1}{v_2}$$

Thus, the refractive index of one medium with respect to another medium is also given by the ratio of the speed of light in first medium to its speed in the second medium.

REFRACTIVE INDEX- NUMERICALS

PROBLEM 4

Speed of light in air is $3 \times 10^8 \text{ m s}^{-1}$ and the speed of light in a medium is $2 \times 10^8 \text{ ms}^{-1}$. Find the refractive index of the medium with respect to air.

Solution:

$$\text{Refractive index } (\mu) = \frac{\text{Speed of light in air } (c)}{\text{Speed of light in the medium } (v)}$$

$$\mu = \frac{3 \times 10^8}{2 \times 10^8} = 1.5$$

PROBLEM 5

Refractive index of water is $4/3$ and the refractive index of glass is $3/2$. Find the refractive index of glass with respect to the refractive index of water.

Solution:

$${}_w\mu_g = \frac{\text{Refractive index of glass}}{\text{Refractive index of water}} = \frac{\frac{3}{2}}{\frac{4}{3}} = \frac{9}{8} = 1.125$$

SNELL'S LAW OF REFRACTION

- Refraction of light rays, as they travel from one medium to another medium, obeys two laws, which are known as Snell's laws of refraction.
- They are: I) The incident ray, the refracted ray and the normal at the point of intersection, all lie in the same plane.
- II) The ratio of the sine of the angle of incidence (i) to the sine of the angle of refraction (r) is equal to the refractive index of the medium, which is a constant.
- $\sin i / \sin r = \mu$

SNELL'S LAW OF REFRACTION




Figure 3.12 Snell's Law

DISPERSION

Activity 7

- Place a prism on a table and keep a white screen near it. Now, with the help of a torch, allow white light to pass through the prism.
- What do you see?
- You can observe that white light splits into seven colored light rays namely, violet, indigo, blue, green, yellow, orange and red (VIBGYOR) on the screen.
- Now, place another prism in its inverted position, between the first prism and the screen.
- Now, what do you observe on the screen?
- You can observe that white light is coming out of the second prism.

DISPERSION

- In the above activity, you can see that the first prism splits the white light into seven coloured light rays and the second prism recombines them into white light, again. Thus, it is clear that white light consists of seven colours. You can also recall the Newton's disc experiment, which you studied in VII standard.



DISPERSION

- Splitting of white light into its seven constituent colours (wavelength), on passing through a transparent medium is known as dispersion of light.
- Why does dispersion occur?
- It is because, light of different colours present in white light have different wavelength and they travel at different speeds in a medium.
- You know that refraction of a light ray in a medium depends on its speed.
-

DISPERSION

- As each coloured light has a different speed, the constituent coloured lights are refracted at different extents, inside the prism.
- Moreover, refraction of a light ray is inversely proportional to its wavelength.
- Thus, the red coloured light, which has a large wavelength, is deviated less while the violet coloured light, which has a short wavelength, is deviated more.

FORMATION OF RAINBOW



The formation of rainbow is an example of dispersion of white light. This can be seen on the opposite side of the Sun. After a rainfall, large number of droplets still remain suspended in the air. When white light passes through them, it is split into seven colours. Dispersion of white light from a large number of droplets eventually forms a rainbow.

